

ize on a UV lightbox. Compare the spotted sample of unknown concentration with the standards to determine the concentration of the cDNA.

8. Since the cDNA is heavily methylated, introduction into a host with an McrA, McrCB, hsdSMR, Mrr phenotype would be subject to digestion by these restriction systems. Therefore, the choice of packaging extract and an *E. coli* host strain is crucial (24–29).
9. Since λ phage can adhere to dead as well as to viable cells, the lower temperature prevents the bacteria from overgrowing.
10. Most cDNA vectors have color selection by Isopropyl- β -D-thio-galactopyranoside (IPTG) and 5-bromo-4-chloro-3-indoyl- β -D-galactopyranoside (X-gal). These components can be added to the top agarose before plating to produce the background blue color. Use 15 μ L of 0.5M IPTG (in water) and 50 μ L of X-gal at 250 mg/mL in dimethylformamide.

References

1. Chomczynski, P. and Sacchi, N. (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* **162**, 156–159.
2. Sambrook, J., Fritsch, E. F., and Maniatis, T., eds. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
3. Han, J. H. and Rutter, W. J. (1987) Lambda gt22, an improved lambda vector for the directional cloning of full-length cDNA. *Nucleic Acids Res.* **15**, 6304.
4. Huynh, T. V., Young, R. A., and Davis, R. W. (1985) *DNA Cloning*, vol. I (Glover, D. A., ed.), IRL, Oxford, Washington, D.C., pp. 49–78.
5. Meissner, P. S., Sisk, W. P., and Berman, M. L. (1987) Bacteriophage lambda cloning system for the construction of directional cDNA libraries. *Proc. Natl. Acad. Sci. USA* **84**, 438–447.
6. Murphy, A. J. M. and Efstratiadis, A. (1987) Cloning vectors for expression of cDNA libraries in mammalian cells. *Proc. Natl. Acad. Sci. USA* **84**, 8277–8281.
7. Palazzolo, M. J. and Meyerowitz, E. M. (1987) A family of lambda phage cDNA cloning vectors, lambda SWAJ, allowing the amplification of RNA sequences. *Gene* **52**, 197–206.
8. Scherer, G., Telford, J., Baldari, C., and Pirrotta, V. (1981) Isolation of cloned genes differentially expressed at early and late stages of Drosophila embryonic development. *Dev. Biol.* **86**, 438–447.
9. Young, R. A. and Davis, R. W. (1983) Efficient isolation of genes by using antibody probes. *Proc. Natl. Acad. Sci. USA* **80**, 1194–1198.
10. Young, R. A. and Davis, R. W. (1983) Yeast RNA polymerase II genes: isolation with antibody probes. *Science* **222**, 778–782.
11. Swaroop, A. and Weissman, S. M. (1988) Charon BS (+) and (–), versatile lambda phage vectors for constructing directional cDNA libraries and their efficient transfer to plasmids. *Nucleic Acids Res.* **16**, 8739.
12. Palazzolo, M. F., Hamilton, B. A., Ding, D. L., Martin, C. H., Mead, D. H., Mierendorf, R. C., Raghavan, K. V., Meyerowitz, E. M., and Lipshitz, H. D. (1990)